

ON MICHELL-LAPLACE DARK BODY

ANGELO LOINGER

ABSTRACT. The dark body of Michell-Laplace has nothing to do with the relativistic black hole.

Many physicists believe that the notion of black hole (BH) is the natural relativistic generalization of the Newtonian notion of dark body (DB), investigated by Michell (1784) and by Laplace (1796).

Now, as it has been shown by McVittie [1] (see also Eisenstaedt [2]), the above belief is based only on “a play of words in expressions such as *the velocity of escape* or *the escape from a body*”. Since a recent article by Stephani [3] has given a further support to this conceptual confusion, it is useful to recall here the very simple argument by McVittie.

Let us indeed consider a celestial spherical body of radius R and mass M . According to Newtonian dynamics, the velocity of escape w of a particle, which is projected radially outwards from the surface of the body, is given by

$$(1) \quad w^2 = 2GM/R,$$

where G is the gravitational constant.

If the particle is projected with a velocity u smaller than w , i.e. if

$$(2) \quad u^2 < 2GM/R,$$

it will arrive at a finite distance from the celestial body, and then it will fall on its surface again.

By using the Newtonian corpuscular theory of light, which says that light is composed of corpuscles obeying Newton’s law of gravitation, and travelling with a given velocity c , Michell and Laplace remarked that if $c < w$ the light corpuscles cannot go away indefinitely from the celestial body. Only if the radius R of the body is such that

$$(3) \quad R = 2GM/c^2,$$

they can escape from the gravitational attraction exerted by the mass M . Then, if

$$(4) \quad R < 2GM/c^2,$$

the light corpuscles will attain a finite distance d from the celestial body, and an observer situated at an intermediate distance between R and d will see the celestial body, owing to the light corpuscles which arrive at his eyes.

If only the spherically symmetrical BH of GR existed – in reality this notion, and the notion of Kerr’s BH, are the product of a misinterpretation of the formalism of GR [4]–, it ought to have the fundamental property that neither the material particles nor the light corpuscles can leave its surface. Therefore, such an object would be invisible to any observer, however near he might be. None of the phenomena observed by the experimentalists in the region surrounding a DB of Michell-Laplace is present in the neighborhood of a BH of GR.

The imagined connection with the Newtonian formula (3) comes out in this way: if for the determination of the Einsteinian gravitational field generated by a point mass M (Schwarzschild’s problem) one chooses the **standard** form – as Droste, Hilbert and Weyl (**but not Schwarzschild** [5]) did – the radial coordinate r_0 of the points of the space surface $r = r_0$ (r_0 is the “radius of the BH”) is given by

$$(5) \quad r_0 \equiv 2GM/c^2;$$

this formula resembles the Newtonian formula (3), which concerns an escape velocity c . But eq.(3) implies obviously that all observers – including those at an infinitely great distance from the celestial body – can see it. As it is clear, the Newtonian dark body of Michell-Laplace is **not** a black hole!

SCHÜLER:

Kann Euch nicht eben ganz verstehen.

MEPHISTOFELES:

Das wird nächstes schon besser gehen,

Wenn Ihr lernt alles reduzieren

Und gehörig klassifizieren.

Goethe, *Faust* (vv.1943-46)

REFERENCES

- [1] G.C. McVittie, *The Observatory*, **78** (1978) 272.
- [2] J. Eisenstaedt, *Gen. Rel. Grav.*, **23** (1991) 75.
- [3] H. Stephani, *arXiv:gr-qc/0304087 v1* (23 April 2003).
- [4] A. Loinger, *Spacetime & Substance*, **3** (2002) 145; Idem, *On Black Holes and Gravitational Waves*, (La Goliardica Pavese, Pavia) 2002, Epitome and Part I.
- [5] K. Schwarzschild, *Berl.Ber.*, (1916) 189; an English translation in *arXiv:physics/9905030* (12 May 1999).

DIPARTIMENTO DI FISICA, UNIVERSITÀ DI MILANO, VIA CELORIA, 16 - 20133 MILANO (ITALY)

E-mail address: `angelo.loinger@mi.infn.it`